

UNITED STATES PLANT PATENT APPLICATION

of

L. PERNILLE AND MOGENS N. OLESEN

for

ROSE PLANT NAMED

'POULpm004'

SUMMARY OF THE INVENTION

BOTANICAL CLASSIFICATION

Rosa hybrida

5

VARIETY DENOMINATION

'POULpm004'

10 The present invention constitutes a new and
distinct variety of garden rose plant which
originated from a controlled crossing between the
female seed parent, an unnamed seedling, and the
male pollen parent, an unnamed seedling. Both
parent plants are non-patented. The two parents
were crossed during the summer of 1997 and the
15 resulting seeds were planted in a controlled
environment in Fredensborg, Denmark. The new variety
is named 'POULpm004'.

20 The new variety may be distinguished from its
unnamed female seed parent by the following
combination of characteristics:

1. While the unnamed seed parent has an
open flower diameter of 120-140 mm;
the same of 'POULpm004' is 80-100 mm.
2. While the petal count of the female
parent is very double, the same of
25 'POULpm004' is double.

The new variety may be distinguished from its

unnamed male pollen parent, by the following
combination of characteristics:

1. While the unnamed pollen parent has a
flower diameter of approximately 60
5 mm; the same of 'POULpm004' is 80-100
mm.
2. While the unnamed pollen parent has a
flower bud length of 45 mm; the same
of 'POULpm004' is 30 mm.
- 10 3. While the unnamed pollen parent has a
pointed ovoid bud form; the same of
'POULpm004' is a globular bud form.

The objective of the hybridization of this rose
variety was to create a new and distinct variety for
15 garden use with unique qualities, such as:

1. Uniform and abundant mauve flowers;
2. Vigorous, but compact growth when
propagated both as a budded rose and on its
own roots;
- 20 3. Exceptional rose scent;
4. Disease resistance.

This combination of qualities is not present in
previously available commercial cultivars of this
type, known to the inventors, and distinguish
25 'POULpm004' from all other varieties of which we are
aware.

As part of their rose development program, L. Pernille Olesen and Mogens N. Olesen germinated the seeds from the aforementioned hybridization in January of 1998 and conducted evaluations on the resulting seedlings in a controlled environment in Fredensborg, Denmark.

'POULpm004' was selected in the spring of 1998 by the inventors as a single plant from the progeny of the aforementioned hybridization.

Asexual reproduction of 'POULpm004' by traditional budding and rooted cuttings was first done by L. Pernille and Mogens N. Olesen in their nursery in Fredensborg, Denmark in July, 1998. This initial and other subsequent asexual propagations conducted in controlled environments have demonstrated that the phenotypical characteristics of 'POULpm004' are true to type and are transmitted from one generation to the next.

20

BRIEF DESCRIPTION OF THE DRAWING

The accompanying color illustration shows as true as is reasonably possible to obtain in color photographs of this type, characteristics of the flower buds, open flowers, leaves, and stems, of 'POULpm004'. Specifically illustrated in SHEET 1:

Fig. 1.1; Stem showing branching, the attachment of leaves, buds, and peduncles, as well as open flowers at various stages of development;

5 Fig. 1.2; Sepals, receptacle, and peduncle.

Specifically illustrated in SHEET 2:

Fig. 2.1; Flower petals and petaloids, detached;

10 Fig. 2.2; Mature leaf, upper surface, as well as juvenile leaf;

Fig. 2.3; Bare stems exhibiting thorns.

DETAILED DESCRIPTION OF THE VARIETY

15 The following is a description of 'POULpm004', as observed in its growth in in a field nursery in Jackson County, Oregon. Observed plants are 3 years of age, and were grown on *Rosa multiflora* understock. Color references are made using the
20 Royal Horticultural Society (London, England) Colour Chart, 1995, except where common terms of color are used.

For a comparison, several physical characteristics of the rose variety 'POULen005', a
25 rose variety from the same inventors described and illustrated in U.S. Plant Patent Application No.

10/738,156 and dated December 16, 2003, are compared to 'POULpm004' in Chart 1.

CHART 1

5		'POULpm004'	'POULen005'
	General tonality.	White Group 155C with intonations of Purple Group 75C - 75D.	Violet Group 84C.
	Diameter of open flower.	75 mm.	80 to 100 mm
10	Petal size.	62 mm (l) x 50 mm (w).	60 to 65 mm (l) x 36 mm (w).

FLOWER AND FLOWER BUD

Blooming habit: Recurrent.

15 Flower bud:

Size: Upon opening, 30 mm in length from base of receptacle to end of bud.
Bud diameter is 23 mm.

20 Bud form: Globular.

Bud color: As sepals unfold, petals are Red Group 39A with intonations of Purple Group Purple Group 79C and Purple

Group 76A. At $\frac{1}{4}$ opening
petals are Violet Group 84C
with intonations of White
Group 155B.

5 Sepals:

Upper Surface:

Upper Surface is Yellow-
Green Group 145C in color.
Moderately pubescent.

10 Lower Surface:

Lower Surface is Yellow-
Green Group 144A. Light
anthocyanic pigments the
color of Greyed-Orange Group
169A-171A observed. Texture
is smooth with scant
stipitate glands.

15

Shape: Sepal apex is cirrhose.
Base is flat at union with
receptacle.

20

Sepal Margin:

Margins have medium
foliaceous appendages on
three of the five sepals.

25

Size: 26 mm long by 11 mm wide.

Receptacle:

	Surface Texture:	Smooth
	Shape:	Funnel shaped.
	Size:	6 mm in length by 5 mm in width.
5	Color:	Yellow-Green Group 144B.
	Peduncle:	
	Surface:	Smooth and glabrous.
10	Length:	53 to 60 mm average length.
	Color:	Yellow-Green Group 144B.
	Strength:	Strong.
15	Borne:	Multiples of 7 buds per flowering stem.
	Flower bloom:	
	Fragrance:	Strong perfume scent.
20	Duration:	The blooms have a duration on the plant of approximately 10 to 14 days. Petals fall cleanly away from plant.
25	Size:	Flower diameter is 75 mm when open. Flower

depth is 45 to 50 mm.

Form: Globular.

Side view flower form:

5 Upon opening;

The upper portion is
flat. The lower
portion is concave.

Open flowers;

10 The upper portion is
flat. The lower portion
is concave.

Petalage: 97 petals on average
under normal conditions
15 with 23 petaloids.

Color:

Upon opening, petals:

Outermost petals:

20 Outer side: White Group 155B, with
intonations of Purple
Group 75C-75D at
marginal zone.

Inner Side: White Group 155C.

Innermost petals:

25 Outer side: White Group 155B with
light intonations of

Purple Group 75C-75D at
marginal zone.

Inner Side: White Group 155C.

5 Upon opening, basal petal spots: No distinctive
coloration at the petal base observed.

After opening, petals:

Outermost petals:

10 Outer side: White Group 155B with
intonations of Purple
Group 75C-75D at
marginal zone.

Inner Side: White Group 155C.

15 Innermost petals:

Outer side: White Group 155B with
intonations of Purple
Group 75C-75D at
marginal zone.

20 Inner Side: White Group 155C.

After opening, basal petal spots: No
distinctive coloration at the petal base
observed.

25

General Tonality: On open flower White Group

5

155C with intonations of
Purple Group 75C-75D. No
change in the general
tonality at the end of the
10th day.

Petals:

10

Petal Reflex: Strongly reflexed.
Margin: Entire and uniform.
Shape: Apex is round. Base shape
varying from acute to round.
Size: 62 mm (l) 50 mm (w).
Texture: Smooth.
Thickness: Thick.
Arrangement: Not formal.

15

Petaloids:

20

Quantity: 20 to 30.
Color:
Upper surface: White Group 155C.
Lower surface: White Group 155B
with intonations of
Purple Group 75C-
75D.
Size: 40 mm (l) x 18 mm
(w).

25

Reproductive Organs:

	Pollen:	None observed.
	Anthers:	
5	Size:	3 mm in length.
	Color:	Yellow-Orange Group 21B.
	Quantity:	182 (actual count).
	Filaments:	
10	Color:	Yellow Group 4C.
	Length:	10 mm.
	Pistils:	
	Length:	10 mm.
	Quantity:	132 (actual count).
15	Stigmas:	Level relative to the length of the filaments and the height of the anthers.
	Color:	Yellow-White Group 158A.
20	Styles:	
	Color:	Yellow-Green Group 150D. Other observed intonations are Red Group 46C.
25	Hips:	None Observed in the

field nursery in
Jackson County Oregon.

PLANT

5

Plant growth: Narrow and bushy. When grown as
a budded field grown plant on
Rosa multiflora understock, the
average height of the plant is 60
to 100 cm.

10

Stems:

Color:

Young wood: Yellow-Green Group
144A.

15

Older wood: Yellow-Green Group
146A.

Surface Texture:

Young wood: Smooth.

20

Older wood: Smooth.

Thorns:

Incidence: 8 thorns per 10 cm of
stem.

Size: Average length: 7 mm.

25

Color: Greyed-Orange Group
165B.

Shape: Deeply concave.

Plant foliage: Normal number of
 leaflets on normal
 leaves in middle of the
 5 stem: 5 leaflets.

Compound leaf: 150 mm (l) x 120 mm
 (w).

Color:

Mature Foliage:

10 Upper surface is: Yellow-Green
 Group 147A.

Lower surface is: Yellow-Green
 Group 147B.

Juvenile foliage:

15 Upper surface is: Yellow-Green
 Group 147A to
 146A.

Lower surface is: Yellow-Green
 Group 147B.

20

Plant leaves and leaflets:

Stipules:

Size: 20 mm in length.

Shape: Linear, slightly broad
 25 based with outward
 extending apices.

	Margins:	Finely serrated with few stipitate glands.
	Color:	Yellow-Green Group 144A.
5	Petiole:	
	Length:	43 mm.
	Above:	
	Color:	Yellow-Green Group 144B.
10	Anthocyanin:	None.
	Underneath:	Thorns.
	Rachis:	
	Length:	73 mm.
	Above:	
15	Color:	Yellow-Green Group 144B.
	Underneath:	Thorns.
	Leaflet:	
	Edge:	Finely serrated.
20	Leaflet size:	72 mm (l) x 55 mm (w).
	Shape:	Ovate to round. Base is rounded. Apex is somewhat round to cuspidate.
25	Texture:	Smooth.
	Thickness:	Thick.

Arrangement: Odd pinnate.

Venation: Reticulate.

Glossiness: Very glossy.

5 **Disease resistance:**

Above average resistance to mildew, rust, black spot, and Botrytis under normal growing conditions in Jackson County, Oregon.

10 **Cold Hardiness:**

The variety 'POULpm004' has been found to be cold tolerant to USDA Cold Hardiness Zone 6.